



Croup

Croup causes swelling around vocal cords and airways. Children usually get croup between 3 months and 5 years of age. It is a scary, but not usually dangerous condition. Croup often occurs several days after a child has mild symptoms of a cold.



SIGNS & SYMPTOMS

- A cough that sounds like a seal's bark.
- Hoarseness
- A high-pitched, squeaky voice when breathing in



Symptoms of croup can also be like symptoms of more serious problems. These include:

- **Epiglottitis.** This is a bacterial infection that can cause the back of the throat to swell up. If the throat is blocked, breathing in is very difficult. Severe respiratory distress can result. Signs of epiglottitis are:
 - Drooling
 - Hanging the head down
 - Sticking out the jaw to breathe
 - Fever
- Something can be stuck in your child's windpipe.



CAUSES

Croup is usually caused by a certain virus. Other viruses, allergies, bacteria, and inhaled foreign objects can mimic croup.

TREATMENT

Self-care measures can treat symptoms. Croup usually goes away in 3 to 7 days. It is usually worse at night. Emergency care is needed for severe problems breathing.



TRIAGE QUESTIONS



Does the child have any of these problems?

- Blue color around the lips or fingernails
- Extreme shortness of breath
- Listlessness or severe weakness
- Coughing so much that your child can't take a breath
- Inability to swallow or make sounds normally

NO

YES ➡ GET MEDICAL CARE FAST



Does the child have a hard time breathing and is he or she doing these things?

- Drooling
- Breathing through the mouth and gasping for air.
- Sticking the chin out

NO

YES ➡ GET MEDICAL CARE FAST



Without having trouble breathing, is the child doing all of the following?

- Drooling
- Breathing through the mouth
- Sticking the chin out

NO

YES ➡ SEE DOCTOR



Does the child make a high, whistling sound like a barking seal?

NO

YES ➡ SEE DOCTOR



USE SELF-CARE



FOR MORE INFORMATION:

The American Academy
of Pediatrics
aap.org



SELF-CARE

- Don't panic. You can help your child stay calm if you stay calm. Hold your child to comfort them. The windpipe may open up a little if your child relaxes. Call your child's doctor or get immediate care if you are not sure what to do.
- Try to keep your child calm. Croup symptoms tend to get worse if your child is crying and upset.
- Go into the bathroom with your child and close the door.
 1. Turn on the hot water in the sink and shower. Let the steam fill the room.
 2. Don't put your child in the shower. Sit with your child. (Don't sit on the floor.) Read a book or play a game with your child. This will help pass the time.
 3. Open the window to let cool air in. This helps make more steam.
 4. Stay in the bathroom about 10 to 15 minutes.
- Use a humidifier in your child's room. Use warm, distilled water, not tap water. Clean the humidifier every day. Put a humidifier on your furnace. Change the filter often.
- If warm mist doesn't help, put warm clothing on your child and take them outside to breath the cold night air for a few minutes.
- Give your child a clear liquid. Warm it first. Warm liquids may help loosen the mucus. Give babies under 6 months old water or electrolyte water, such as Pedialyte, if okay with your child's doctor. Give water, diluted apple juice, etc. to a child who is 6 months old or older.

To Help Prevent Croup and Other Infections

- See that your child's immunizations for diphtheria (DTaP), measles (MMR), and H. influenzae type b (Hib) are up to date.
- Follow prevention measures for colds and flu.